

1,3-Dipolar cycloaddition of C-Benzoyl-N-phenyl- and C,N-diphenylnitrones to β -substituted vinylphosphonates

Arbuzov B., Lisin A., Dianova E., Samitov Y.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

1. The 1,3-dipolar cycloaddition of C,N-diphenyl- and C-benzoyl-N-phenylnitrones to diethyl β,β -diacetylvinylphosphonate yields trans-2,3-diphenyl-4-diethoxyphosphono-5,5-diacetylisoaxazolidine and trans-2-phenyl-3-benzoyl-4-diethoxyphosphono-5,5-diacetylisoaxazolidine, respectively. 2. C-Benzoyl-N-phenylnitrone reacts with diethyl β -phenylvinylphosphonate to form trans-trans-2,5-diphenyl-3-benzoyl-4-diethoxyphosphonoisoaxazolidine, and it reacts with dimethyl β -ethoxyvinylphosphonate to form trans-trans-2-phenyl-3-benzoyl-4-dimethoxyphosphono-5-ethoxy-isoaxazolidine. 3. As the electrophilic character of the double bond in the β -substituted vinylphosphonates is enhanced, their reactivity in 1,3-dipolar cycloaddition to nitrones increases. © 1979 Plenum Publishing Corporation.

<http://dx.doi.org/10.1007/BF00946684>
